



W.U.S. HEALTH CENTRE
UNIVERSITY OF DELHI

DENGUE FEVER : ADVISORY

WUSHC/17-18/980

dt 30/6/17

To be forwarded
to all HODs/Deans
colleges
& HCs

Neeb Selgi
4/7/17

Dengue fever is a mosquito-borne tropical disease caused by the dengue virus.

Symptoms and Signs :- Dengue triad: High fever, severe headache, skin rash

1. Sudden onset high fever biphasic or saddleback in nature, breaking & returning.
2. Severe headache behind the eyes, severe muscle and joint pains.
3. Characteristic skin rash similar to measles. In some it develops into the life-threatening dengue hemorrhagic fever, resulting in bleeding, low levels of platelets and blood plasma leakage or into dengue shock syndrome when low blood pressure occurs.
4. During critical phase : lymphadenopathy, mouth & nose bleeding, low blood pressure, accumulation of fluid in the chest, accumulation of fluid in abdominal cavity, depletion of fluid, organ dysfunction, gastrointestinal bleeding.
5. During recovery phase : altered level of consciousness, seizures, itching, slow heart rate, peeling of the skin, slow heart rate, fatigue.
6. Dengue shock syndrome.
7. Dengue hemorrhagic fever – usually affects children less than 10 years of age.

The incubation period ranges from 4 to 7 days.

Diagnosis :-

On examination :- Lymphadenopathy, Pleural effusions, Ascites.

- Low white blood cell count (WBC), positive tourniquet test or any warning sign.
- The earliest change is a low white blood cell count, low platelets and metabolic acidosis. Elevated level of Aminotransferase (AST and ALT) rising hematocrit and hypoalbuminemia.
- Check for the virus or antibodies to the virus.


Treatment :-

- No specific treatment. There is no vaccine to prevent dengue fever.
- Drink plenty of fluids.
- Intravenous fluids with electrolyte replacement to maintain urinary output of 1 ml/kg/hr.
- Acetaminophen can alleviate pain and reduce fever.
- Avoid Aspirin, Ibuprofen and Naproxen Sodium.
- Transfusion with packed red blood cells or whole blood to replace blood loss.
- During recovery phase intravenous fluids are discontinued to prevent a state of fluid overload or else loop diuretic e.g. furosemide if the patient is outside the critical phase.

Prevention :-

- Stay away from heavily populated residential areas, if possible.
- Use mosquito repellents, even while indoors.
- When indoors, live in air conditioned room and in well screened houses.
- Make sure window and door screens are secure and free of holes. If sleeping areas are not screened properly or air conditioned, use mosquito nets.
- To decrease mosquito population get rid of old automobile tires, cans, flower pots.
- Use insect repellent 10% DEET, Permethrin.
- Generalised spraying of environs with Organophosphate or Pyrethroid Insecticides.
- Overhead water tank (OHT)/cemented tanks should have well fitted lid and should be kept locked.
- Water outlet/air bent pipe of OHT should be covered with metallic/plastic net of sufficient size.
- Ensure that there is no water logging at roof tops and in the ground areas.
- Water contained in money plant/bamboo plant should be changed at least once a week.
- Water contained in coolers should be replaced after scrub cleaning weekly and if not possible to change water, add 2 tablespoon full of petrol/1table spoon temiphos granules.
- Use mesh on doors and windows.
- Wear full sleeves cloths and trousers that cover arm and legs.

30th June, 2017


(Dr. Sunil)
Chief Medical Officer

Copy to : Joint Registrar Secretariat of Hon'ble Vice Chancellor, Pro-Vice Chancellor, Director South Campus, Dean (Colleges), Treasurer, Proctor, Registrar, Chairman – ICH, W.U.S. Health Centres, Finance Officer, Librarian, all Faculties/Departments/Colleges – All the head of Departments are requested to designate a Nodal Officer so that there are no mosquitogenic conditions in their office.

Please disseminate to all and display it on Notice Board

Chief Medical Officer's Office

Diary No. 664
Date 3/7/17